



## The translocation of microplastics to lipid droplets of *Daphnia magna* is an artefact

Schür, Christoph ; Rist, Sinja; Hartmann, Nanna B.; Wagner, Martin

*Published in:*

MICRO 2018 Fate and Impact of Microplastics: Knowledge, Actions and Solutions - conference proceedings

*Publication date:*

2018

*Document Version*

Version created as part of publication process; publisher's layout; not normally made publicly available

[Link back to DTU Orbit](#)

*Citation (APA):*

Schür, C., Rist, S., Hartmann, N. B., & Wagner, M. (2018). The translocation of microplastics to lipid droplets of *Daphnia magna* is an artefact. In J. Baztan, M. Bergmann, A. Carrasco, C. Fossi, B. Jorgensen, A. Miguelez, S. Pahl, R. C. Thompson, & J-P. Vanderlinden (Eds.), *MICRO 2018 Fate and Impact of Microplastics: Knowledge, Actions and Solutions - conference proceedings* (pp. 234-234) <https://micro2018.sciencesconf.org/resource/page/id/8>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# The translocation of nano- and microplastics to lipid droplets of *Daphnia magna* is an artefact

Christoph Schür<sup>1</sup>, Sinja Rist<sup>2</sup>, Nanna Bloch Hartmann<sup>2</sup>, Martin Wagner<sup>3</sup>

<sup>1</sup>Department of Aquatic Ecotoxicology, Goethe University Frankfurt am Main, Max-von-Laue-Str. 13, 60438 Frankfurt am Main, Germany

<sup>2</sup>Department of Environmental Engineering, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

<sup>3</sup>Department of Biology, Norwegian University of Science and Technology, Realfagbygget, DU1-115, Gløshaugen, Norway

The last decade has seen a surge in research investigating various aspects of micro- and nanoplastics, including occurrence, uptake, and potential effects in biota. Our knowledge about biota-particle-interactions is still limited and often based on early studies that – due to the infancy of the field – may have deficiencies in the experimental design and quality controls. One such example relates to the potential of plastic particles to cross the epithelium of the digestive tract and translocate to other tissues. This has been reported in the literature for the freshwater cladoceran *Daphnia magna* and – if true – is toxicologically relevant. The aim of our study was to replicate previous findings while covering additional experimental scenarios.

We exposed 205 starved *Daphnia magna* neonates to two particle types (fluorescent polystyrene spheres of 20 nm and 1000 nm diameter) at two concentrations (2 µg/L and 2 mg/L) for two time periods (4 h and 24 h). Additionally, we adapted a tissue clearing method to the use with *Daphnia* to improve visibility of particles inside the specimen. We used confocal laser scanning microscopy to investigate the translocation of particles from the gut to other tissues, including lipid storage droplets. This is supplemented by experiments regarding the leaching of the dye from the particles.

The 1000 nm particles can be imaged individually inside the specimen. At the higher particle concentration we detected fluorescence in the lipid droplets that for the 1000 nm particles was clearly separate from particles and faded within minutes of observation. Our findings indicate that the tissue translocation to the lipid storage droplets reported previously is probably an experimental artefact caused by the leaching of the dye used in commercially available plastic beads. This underlines the importance of scepticism and replication in our field, especially if a plausible underlying biological mechanism is lacking.